

## Integrating climate change criteria in reforestation projects using a hybrid decision-support system

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 Environ. Res. Lett. 10 094022

(<http://iopscience.iop.org/1748-9326/10/9/094022>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.105

This content was downloaded on 13/04/2017 at 08:22

Please note that [terms and conditions apply](#).

You may also be interested in:

[Spatial Analytic Hierarchy Process Model for Flood Forecasting: An Integrated Approach](#)

Abd Nasir Matori, Dano Umar Lawal, Khamaruzaman Wan Yusof et al.

[The Importance of Site Selection for Radio Astronomy](#)

Roslan Umar, Zamri Zainal Abidin and Zainol Abidin Ibrahim

[How countries link REDD+ interventions to drivers in their readiness plans: implications for monitoring systems](#)

G Salvini, M Herold, V De Sy et al.

[Forests in a water limited world under climate change](#)

Csaba Mátyás and Ge Sun

[Operational approaches to managing forests of the future in Mediterranean regions within a context of changing climates](#)

Scott L Stephens, Constance I Millar and Brandon M Collins

[Modeling groundwater vulnerability to pollution using Optimized DRASTIC model](#)

Kehinde Anthony Mogaji, Hwee San Lim and Khiruddin Abdullar

[Measurement and monitoring needs, capabilities and potential for addressing reduced emissions from deforestation and forest degradation under REDD+](#)

Scott J Goetz, Matthew Hansen, Richard A Houghton et al.

[Making environmental assessments of biomass production systems comparable worldwide](#)

Markus A Meyer, Ralf Seppelt, Felix Witing et al.

## Environmental Research Letters



## LETTER

## OPEN ACCESS

RECEIVED  
27 April 2015

REVISED  
27 August 2015

ACCEPTED FOR PUBLICATION  
27 August 2015

PUBLISHED  
16 September 2015

Content from this work  
may be used under the  
terms of the [Creative  
Commons Attribution 3.0  
licence](#).

Any further distribution of  
this work must maintain  
attribution to the  
author(s) and the title of  
the work, journal citation  
and DOI.



# Integrating climate change criteria in reforestation projects using a hybrid decision-support system

Jorge Curiel-Esparza<sup>1</sup>, Nuria Gonzalez-Utrillas<sup>2</sup>, Julian Canto-Perello<sup>3,4</sup> and Manuel Martin-Utrillas<sup>1</sup>

<sup>1</sup> Physical Technologies Center, Universitat Politecnica de Valencia, Camino de Vera s/n, E-46022 Valencia, Spain

<sup>2</sup> Doctoral School, Universitat Politecnica de Valencia, Camino de Vera s/n, E-46022 Valencia, Spain

<sup>3</sup> Department of Construction Engineering and Civil Engineering Projects, Universitat Politecnica de Valencia, Camino de Vera s/n, E-46022 Valencia, Spain

<sup>4</sup> Author to whom correspondence should be addressed

E-mail: [jcuriel@fis.upv.es](mailto:jcuriel@fis.upv.es), [mgmartin@fis.upv.es](mailto:mgmartin@fis.upv.es), [nugonut@upv.es](mailto:nugonut@upv.es) and [jcantope@cst.upv.es](mailto:jcantope@cst.upv.es)

**Keywords:** AHP, Delphi, reforestation, climate change, Mediterranean region

## Abstract

The selection of appropriate species in a reforestation project has always been a complex decision-making problem in which, due mostly to government policies and other stakeholders, not only economic criteria but also other environmental issues interact. Climate change has not usually been taken into account in traditional reforestation decision-making strategies and management procedures. Moreover, there is a lack of agreement on the percentage of each one of the species in reforestation planning, which is usually calculated in a discretionary way. In this context, an effective multicriteria technique has been developed in order to improve the process of selecting species for reforestation in the Mediterranean region of Spain. A hybrid Delphi-AHP methodology is proposed, which includes a consistency analysis in order to reduce random choices. As a result, this technique provides an optimal percentage distribution of the appropriate species to be used in reforestation planning. The highest values of the weight given for each subcriteria corresponded to FR (fire forest response) and PR (pests and diseases risk), because of the increasing importance of the impact of climate change in the forest. However, CB (conservation of biodiversity) was in the third position in line with the aim of reforestation. Therefore, the most suitable species were *Quercus faginea* (19.75%) and *Quercus ilex* (19.35%), which offer a good balance between all the factors affecting the success and viability of reforestation.

## 1. Introduction

On 24th September 2008, the current Secretary-General of the United Nations stated with reference to the United Nations Collaborative Program on Reducing Emissions from Deforestation and Forest Degradation in Developing Countries (UN-REDD) that climate change cannot be mitigated without the proactive management of the world's forests. This, however, will be a complex and challenging feat (Kimoon 2008). Monitoring the effectiveness of planned interventions is a key factor in developing reforestation projects (Salvini *et al* 2014). Following this line, a new European Union Forest Strategy for forests and the forest-based sector has been proposed (EUCOM 2013), implementing the strategy adopted

(EUCON 1999) and related to the EU Forest Action Plan (EUCOM 2006). The EU Forest Strategy (EUCOM 1998) points out the challenges facing the policy and legal framework for forests in the EU as well as the common forest policy objectives for EU Member States. This European strategy emphasizes the importance of the forests' multifunctional role in the development of society. It shows that forests and forestry can successfully provide society with multiple benefits. Thus the European Parliament (EUPAR 2011) considers that Mediterranean forests are highly valued sinks of atmospheric carbon and sources of European diversity and should therefore benefit from enhanced protection. Furthermore, the EU biodiversity strategy to 2020 (EUCOM 2011) encourages Member States to ensure that forestation is

carried out in accordance with the pan-European operational level guidelines for sustainable forest management. In order to develop these EU targets and to reduce the alarmingly high rate of global deforestation, reforestation, together with afforestation, forest rehabilitation and other means of rehabilitation, play an important role in EU forest policy.

According to the EU Forest Strategy, European forests serve different purposes such as social (contribution to rural development, recreation), economical (timber, cellulose, packaging, paper, renewable energy), environmental (conservation of biological diversity, action to combat desertification, carbon sequestration which mitigates climate change, protection against soil erosion, avalanche control, regulation and enhancement of streams and rivers) and societal (recreation, employment in rural areas). Guidelines and recommendations from international organizations such as FAO (FAO 2013), IPCC (IPCC 2014a and 2014b), CPF (CPF 2008) and IUNC (IUNC 2004) could also be taken into account. Most of those documents incorporate specific sections for each of the six continents, so a wide range of indicators can be extracted from them according to different environmental conditions worldwide. Hence forest systems are inherently complex and there are several criteria that may be considered in the evaluation of the different species for a reforestation project. Moreover, the specific features of Mediterranean forests highlight the need for achieving a successful reforestation, as the selection of appropriate species is a vital component in a reforestation project, because of the significant long-term implications of such a decision for management costs and the value to the final community. Moreover, the complexity is currently increasing because of the way in which different social groups or stakeholders perceive the relative importance of the criteria involved in the decision-making process (Diaz-Balteiro and Romero 2008, Akiner and Akiner 2010, Hunt *et al* 2014). In view of these difficulties, today's reforestation planning calls for the use of expert systems as a versatile tool for decision support (Orsi *et al* 2011). In this context, the regional forest action plan for Comunitat Valenciana (Spain) forces decision makers to use multicriteria techniques in order to select appropriate species in reforestation planning (PATFOR 2013).

Under Mediterranean climatic conditions, the main problems are drought (Sanchez-Salguero *et al* 2012) and forest fires (Amraoui *et al* 2015), and related to that or as a consequence of them can be mentioned a long time sunlight exposure (Kaya and Kahraman 2011), soil erosion (Ramos and Martinez-Casasnovas 2015) and pest outbreaks (Hódar *et al* 2012). Different criteria might be chosen by the panel of experts, such as ozone concentrations (Paoletti 2005). In fact, in different latitudes may pose a problem just the contrary, the lack of sunlight (Hitsuma *et al* 2015) or waterlogged soils after a flood (Velmurugan *et al* 2015), indeed any other issues such as

snow-induced damage on branches (Kane and Finn 2014), the lack of soil nutrients (Sullivan *et al* 2015) like permafrost affected soils (Dymov *et al* 2015) or the presence of pollutants in soils (Hockmann *et al* 2015). There are other criteria related to the aim of the forestation, such as biodiversity conservation, tourist attractiveness (TA) and wildlife and landscape improvement. They are general enough criteria that can be applied to different aims, such as environmental protection or restoration. However, in the specific case of timber production, different criteria should be taken into account, such as incomes or revenues, log properties such as strengthening or flexure or method of forestation, among others (Denzler *et al* 2015, Hampel *et al* 2015) regarding economic criteria, they are general enough and can be applied to any study case. Nevertheless, other criteria can be added depending the specific features of the forestation project.

This study proposes a decision-support system integrating climate change criteria for optimal reforestation planning using the Delphi Method and Analytical Hierarchy Process (AHP). The methodology is applied to a case study of the Spanish forest in the Mediterranean region. Delphi methodology is used to assess the suitability of various species compatible with the ecological features of the reforesting stand, when considering for instance climatic, physiographical and edaphic factors and potential vegetation. Such species are evaluated according to the purpose of reforestation with reference to several defined criteria proposed by the planning strategies. The task involves determining a ranking of the criteria and their priority using AHP methodology (Saaty 2013, Martin-Utrillas *et al* 2015a). The proposed innovation includes the ecological and socio-economic repercussions of climate change, which are not taken into account in traditional decision-making strategies, in order to reduce the adverse effects of climate change. As a result, the combined Delphi-AHP methodology provides an optimum percentage distribution of the appropriate species to be used in the planning of reforestation incorporating adaptation to climate change.

## 2. Applied methodology for selecting species

Species able to respond to climate change do so by distributional or phenological shifts, acclimating or adapting (Easterling *et al* 2000, Parmesan 2006, Chen *et al* 2011). Newman *et al* (2011) focuses on the processes by which species respond to climate change, whereas Hannah (2010) offers a more applied perspective. Current researches' (Pettorelli 2012) aim to address the underlying mechanisms of species response and forecast species interactions. Extensive phenological data sets, including those presented in Fitter and Fitterm (2002) and Lenoir *et al* (2008), reveal

how the characteristics of species influence the magnitude of phenological shifts. And these shifts may indicate effective acclimation and viability of species. Sexton *et al* (2009) reviews the range limits crucial to species responses to environmental change. Hofmann and Todgham (2010) reviews physiological mechanisms by which organisms may respond to climate change and Gilman *et al* (2010) proposes that identifying the key interactions provides a way to predict how biotic interactions influence responses to climate change. According to this model, DeLucia *et al* (2012) points to the importance of factors such as biotic interactions between plants and insects and Lau and Lennon (2012) provides an example of how the evolution of interacting plant species with rapid life cycles and soil microbial communities may influence responses to environmental changes like drought stress. Finally, Williams *et al* (2008) proposes a framework for assessing the vulnerability of species to climate change, including biotic vulnerability, exposure to climate change, potential evolutionary and acclimatory responses and the potential efficacy of management strategies. As a conclusion, Bucley (2014) highlights the need to integrate accurately forecast species responses in many traditional fields such as physiology, evolution, population and community ecology.

Species suitable for reforestation are determined by using the Delphi technique. Marques *et al* (2013) apply a Delphi survey aimed at identifying empirical guidelines to assist developers and users of decision-support systems for forest management. Experts are given a preliminary list of possibly suitable species according to bioclimatic zone (Gomez-Aparicio *et al* 2011), phytoclimatic versatility (Garcia-Lopez and Allue-Camacho 2008), ecophysiology (Cabrera 2002), physiographical variables (Boucher *et al* 2014) and stand conditions such as soil features, orientation and altitude (Urli *et al* 2014). The best alternatives for the area must be identified, bearing in mind the main goal of reforestation, i.e. biodiversity conservation. Orsi *et al* (2011) performed a Delphi process aimed at defining the key ecological criteria and indicators when considering biodiversity conservation as the main objective of reforestation. At this first stage, a Delphi methodology provides a significant basis for arriving at a consensus.

Delphi methodology is based on expert surveys in several rounds in which the survey results from the previous round are given as feedback on the second and subsequent rounds. The objective of this technique is to obtain the most reliable consensus of opinion from a group of experts (Garson 2013). As a result of this procedure, judgments of a panel of experts are elicited and refined (Meddour-Sahar *et al* 2013). Therefore, Delphi main features are iteration, feedback and anonymity of responses. The group of experts is made up of staff of nature reserves, regional officers, environmental organizations and forestry researchers. They


are given a list of species to be evaluated on a numerical scale according to their suitability for the region's natural environment.

Once the appropriate species have been determined, they are compared in pairs with each one of the criteria and translated into numerical values. These criteria and subcriteria represent economical aspects, environmental services and effects of climate change on forest biodiversity that are expected to arise and could not be adequately managed by conventional approaches to conservation, as shown in figure 1. AHP technique is used to evaluate criteria according to the objective of biodiversity conservation of reforestation, dealing not only with traditional economic and environmental aspects (Martin-Utrillas *et al* 2015b), but also with climate change factors. There is a need to provide a method which allows policy-makers and stakeholders to take appropriate measures to adapt to climate change. The proposed decision-support system for selecting species combines a Delphi-AHP model and provides a systematic approach to evaluating multi-criteria and multi-species problems.

The AHP methodology provides a framework within which these selected species can be prioritized while giving a specific weight to the different criteria (Saaty 2008). This method allows the decision maker to arrive at judgments with regard to the species and the criteria according to which the species are evaluated. Moreover, there are non-commensurable criteria that should be considered simultaneously in order to arrive at a more accurate selection (Canto-Perello *et al* 2013, Curiel-Esparza *et al* 2014). The target of this second stage of reforestation planning is a prioritized list of species, a measure of their relative priority and an index denoting the consistency of the decision maker. As shown in figure 1, the AHP process is developed using a three-level hierarchical structure: criteria, sub-criteria and species.

### 3. Determining criteria and subcriteria for species selection

The criteria and subcriteria have been chosen by the panel of experts taking into account guidelines and recommendations from: the Climate Change Guidelines for Forest Managers (FAO 2013); the Intergovernmental Panel on Climate Change (IPCC 2014a, 2014b); the Ministerial Conferences on the Protection of Forests in Europe (MCPFE 2002, 2007, 2011); the Strategic Framework for Forests and Climate Change proposed by the Collaborative Partnership on Forests (CPF 2008), and from the International Union for Conservation of Nature's Guidelines for Afforestation and Reforestation for Climate Change Mitigation (IUNC 2004). The first criterion relates to the effect on various environmental services (EES) provided by multifunctional forests. The second criterion is connected with the



economic importance (EIM) of forests. The third criterion is related to the impact of climate change (ICC) on the forest ecosystem (Garcia-Lopez and Allue-Camacho 2010).

For the first criterion, that is, the effect on various environmental services, the following subcriteria are considered:

- Contribution to biodiversity (CB). This refers to the uniqueness of a vegetation type, which cannot be found elsewhere or under different conditions (Cram *et al* 2006). Concepts such as autochthony, gregariousness, species richness and maturity level are considered here. Ruiz-Benito *et al* (2013) claim that differences in species richness between planted forests and natural ecosystems might be reduced through management options, allowing more natural conditions to be restored.
- Landscape enhancement (LE). Chromatic heterogeneity, average height from the upper storey and gap distribution patterns might be included in this concept (Gomontean *et al* 2008). Forest management towards more naturalness and wilderness may enhance the landscape (Lupp *et al* 2013). On the other hand, Moreno *et al* (2011) report that landscape variables such as land-cover type, distances to roads and towns, aspect or elevation might influence forest risk.

- TA may be defined here as the response of society to the condition of the environment (Roura-Pascual *et al* 2009), including social acceptance of the species. The public has a right of access for recreational purposes and of indicating the intensity of use. On the other hand, Edwards *et al* (2012a) have carried out a survey of public preferences for forests as sites for recreational use and define recreational value of a forest in terms of the preference of people who regularly use forests as recreation sites.
- Wildlife improvement (WI) refers to the conditions for setting wildlife species in the area. Dietz and Jorgenson (2014) suggest that high levels of human well-being are associated with low pressure on the environment. Ezebilo (2012) studies forest stakeholder participation in improving wildlife habitat in planning a more sustainable forest strategy which integrates wildlife. Other studies assess the effects of forest management on wildlife (Nielsen and Treue 2012). Moreover, maintenance of other socio-economic functions and conditions of forests is required by the EU Forest Strategy (EUCOM 2013). The latter considers safeguarding the forests' natural habitats and their ecological functions to be an essential role.

In the second criterion, i.e. the economic importance of forests, the subcriteria considered take into account not only general factors mentioned in any



reforestation project, but also those related to the significant economic value arising from the multiple benefits provided by biodiversity. According to the EU Biodiversity Strategy to 2020 (EUCOM 2011), these are seldom seized by markets. In this context, the general recommendations for forestation and reforestation projects of the Ministerial Conference on the Protection of Forests in Europe (MCPFE 2007) encourage their contribution to maintaining or improving the provision of ecosystem goods and services. The proposed subcriteria are:

- Initial cost (IC) includes items such as soil preparation cost, planting cost, vegetation clearing, and other socio-economic factors (Barajas-Guzman and Barradas 2013). It also includes replanting cost. Government incentives, subsidies or any financial compensation are taken into account (Zhou *et al* 2007).
- The subcriterion of silvicultural treatments (ST) includes the number and cost of later silvicultural interventions, such as thinning and harvesting regimes (Edwards *et al* 2012b). These ST have a positive effect on the survival and growth of seedlings (Navarro-Cerrillo *et al* 2011). Silvicultural operations may have either positive or negative effects on biodiversity and water protection without high cost (Koprowski and Dunker 2012). Thus, it is important to anticipate the long-term effects of ST in order to develop a sustainable forest management.
- Economic return (ER) refers to the valuation of both market and non-market services, including environmental externalities (Mendoza and Martins 2006) since forests provide important ecosystem services which supply societal needs. On the one hand, Ojea *et al* (2012) compare the economic returns of a sustainable and a non-sustainable forest by analyzing long term trends, showing that sustainable forests yield higher economic benefits. On the other hand, Xu *et al* (2013) estimate the potential of carbon storage in the live biomass of forest in order to reduce emissions of carbon dioxide by restoring cleared and disturbed spruce forest through afforestation and reforestation programs.

The third criterion, i.e. the impact of climate change on forests, includes subcriteria which increase forest ecosystem vulnerability as a result of climate change. These may be declining soil fertility and water availability and increasing risks of forest fires (Kukavskaya *et al* 2013) especially in the Mediterranean regions (Schröter *et al* 2005). Moreover, due to rising temperatures and hotter and drier summers, wildfires are forecast to increase in frequency, intensity and severity (Vautard *et al* 2014). In fact, evidence of the relationship between drought and forest fires has been

reported by Vazquez *et al* (2002) in Northern Spain, regarding the active hot spots detected in the satellite during the four summer months of the year 2000. Other effects of climate change such as droughts, tree mortality, loss of biodiversity and vegetation stress produce increasing fuel loading, insect outbreaks and the spread of invasive species (Vicente-Serrano *et al* 2014). In IPCC 2014a, 2014b Working Group II Fifth Assessment Report, vulnerability is defined as the propensity or predisposition to be adversely affected, which includes concepts such as sensitivity or susceptibility to harm and lack of adaptation to those effects produced by climate change. In the context of forests, impacts attributed to climate change are: increasing burnt forest areas, wildfire frequency and duration; regional increases in tree mortality and insect infestations in forests; rainforest degradation and recession. Reforestation is considered, among other actions, as a policy of adaptation to these impacts, otherwise forest dieback involves risk for carbon storage, biodiversity, wood production, water quality, amenity and economic activity. This adaptation can only occur through policies taking into account rural decision-making contexts which values non-marketed ecosystem services. Trees may be adapted to future climate changes throughout their phenotypic plasticity (Benito Garzon *et al* 2011). Other attributes of species, such as ecological resistance (capacity to maintain integrity under stress) and resilience (capacity to reestablish after disturbance) can attenuate the effects of climate change. Thus, the following subcriteria have been taken into account:

- Water stress response (WR). In the west Mediterranean basin, water stress is considered the most limiting factor for ecosystem reconstruction and the main factor of forest decline (Sanchez-Salguero *et al* 2012). The effects of climate change are rising temperatures, decreasing rainfall and concentration of rainfall in extreme events. Urbietta *et al* (2008) suggest that tree species can differ in their responses to water availability during the phase of establishment.
- Sunlight tolerance (SU). Species tolerate and require a specific amount and period of sunlight at an early age in order to continue their successful development (Peman Garcia *et al* 2006). Sanchez-Gomez *et al* (2006) reported the survival of seedlings of different species under experimental gradients of irradiance and water availability. On the other hand, under sunny conditions, evaporation and transpiration are increased and less water is retained in the soil (Kaya and Kahraman 2011).
- Soil conservation (SC). Reforestation contributes to the prevention of soil erosion since trees intercept rain and slow down the water hitting the soil, which is the umbrella function. Moreover, the soil anchor

function of the roots prevents the soil from washing away (Kaya and Kahraman 2011). Consequently, these functions contribute to soil erosion prevention. Eventually, forest fire can affect soil properties such as aggregate stability, organic matter content, soil microbiology, water repellency and soil mineralogy (Mataix-Solera et al 2011). Selkimaki et al (2012) studied the parameters of forests that are related to surface erosion, associated with climatic conditions, tree species composition and stand structure.

- Forest fire response (FR). Forest fires affect the production of water in the watersheds (Kinoshita and Hogue 2015). Some non-serotinous species, such as *Pinus nigra*, have very low or nil post-fire regeneration (Espelta et al 2003), which is taken into account in the framework of assessing the best alternatives. Moreover, the importance of fire in stimulating population growth of some serotinous species (*Acacia* sp., *Pinus halepensis*, *Pinus pinaster*) and its effect on stimulating germination of soil-stored seeds are important factors when assessing alternatives (Roura Pascual et al 2009). Gonzalez and Pukkala (2007) analyzed relationships between forest characteristics such as species composition and the probability of fire occurrence.
- Pests and diseases risk (PR). Biotic and abiotic damaging agents are forest disturbances, especially in vegetation with a slow growth rate, long life cycle and long reproductive periods (Cram et al 2006). Besides, the management effects on natural ecosystems may not be immediately apparent, and considerable periods may elapse before the long-term effects of anthropogenic perturbations become clear (Reilly and Elder 2014).

Thus, three criteria can be distinguished in the hierarchical model. EES considers the effect on several environmental services, EIM includes some economic factors and ICC takes into account the impact of the climate change on forests. For each criterion and sub-criterion, species are ranked to obtain an overall preference percentage for them. AHP methodology is required to evaluate these criteria in order to assign a specific priority to each factor and to evaluate the relative importance of the species in terms of each criterion and sub-criterion.

#### 4. Evaluation of expert judgments and pairwise comparison

The process is performed in three stages. In the first step, the three criteria are compared in pairs by the panel of experts using a questionnaire with a 9-point scale (see table 1). Higher values indicate a greater priority and reciprocals are used to show their preference for the inverse choice (Curiel-Esparza and

Canto-Perello 2013, Martin-Utrillas et al 2015c). The panel of experts is requested to assess the three criteria as is shown in table 2. The priorities of each group are established in this way, providing as a result a pairwise comparison matrix of the criteria EES, EIM and ICC:

$$[M] = \begin{bmatrix} 1.0000 & 0.9583 & 0.3476 \\ 1.0435 & 1.0000 & 0.2231 \\ 2.8769 & 4.4815 & 1.0000 \end{bmatrix}. \quad (1)$$

Thereafter, the priority vector constituted by the percentages of overall relative priorities can be obtained by the eigenvector method:

$$[\omega] = \begin{bmatrix} 0.1897 \\ 0.1684 \\ 0.6419 \end{bmatrix}. \quad (2)$$

In order to evaluate how coherent the panel of experts is with respect to their answers, a consistency analysis is developed calculating the eigenvalue ( $\lambda_{\max} = 3.0263$ ) to obtain the consistency ratio (CR). Then, the consistency index is defined as:

$$CI = \frac{\lambda_{\max} - n}{n - 1} = 0.0131, \quad (3)$$

where  $n$  is the order of the matrix. The random consistency index (RCI) for  $n = 3$  is 0.52 (Saaty 2012). Therefore, the CR is calculated as follows:

$$CR = \frac{CI}{RCI} = 2.53\%. \quad (4)$$

Maximum CR values are given depending on  $n$  value. In this case ( $n = 3$ ), CR must be below 5%, hence the results are reliable.

In the second survey, the panel of experts is requested to compare the twelve subcriteria as shown in table 2. Subcriteria in each group are assessed in order to determine their relative weights, using the eigenvector method (see table 3 and figure 2). Once again, the priorities of each subcriterion within each criterion are established to evaluate their weight. In this case, the pairwise comparison matrix of the four subcriteria in the first criterion (EES) for the panel of experts is as follows:

$$[M_{EES}] = \begin{bmatrix} 1.0000 & 2.9808 & 8.3922 & 6.9880 \\ 0.3355 & 1.0000 & 6.8525 & 4.4922 \\ 0.1192 & 0.1459 & 1.0000 & 0.3704 \\ 0.1431 & 0.2226 & 2.6999 & 1.0000 \end{bmatrix}. \quad (5)$$

The eigenvector method is applied, pointing out the percentages of overall relative priorities for each factor on the EES criterion:

$$[\omega_{EES}] = \begin{bmatrix} 0.5836 \\ 0.2848 \\ 0.0455 \\ 0.0860 \end{bmatrix}. \quad (6)$$

And the parameters of the consistency analysis in this case adopt the following values: 4.1390 for the  $\lambda_{\max}$ , the RCI ( $CI = 0.0463$ ) and the CR ( $CR = 5.21\%$ ), which is below 9% for  $n = 4$  in this case, so it can be

**Table 1.** Saaty's fundamental scale for pairwise comparisons used by the panel of experts in the APH methodology (Saaty 2012).

Intensity of importance	Definition	Explanation
1	Equal importance	Two factors or technologies contribute equally to the goal
3	Moderate importance	Experience and judgment slightly favor one factor or technology over another
5	Strong importance	Experience and judgment strongly favor one factor or technology over another
7	Very strong or demonstrated	A factor or technology is favored very strongly over another
9	Extreme importance	The evidence favoring one factor or technology over another is of the highest possible order of affirmation
1/3 1/5 1/7 1/9	Reciprocals of above	If factor or technology i has one of the above non-zero numbers assigned to it when compared with activity j, then j has the reciprocal value when compared with i

**Table 2.** Questionnaire given to the panel of experts in order to evaluate criteria and subcriteria in a pair-wise comparison, according to the Saaty's fundamental scale shown in table 1.

Main groups	More important than				Equal	Less important than				Main groups
EES	9	7	5	3	1	3	5	7	9	EIM
EES	9	7	5	3	1	3	5	7	9	ICC
EIM	9	7	5	3	1	3	5	7	9	ICC
CB	9	7	5	3	1	3	5	7	9	LE
CB	9	7	5	3	1	3	5	7	9	TA
CB	9	7	5	3	1	3	5	7	9	WI
LE	9	7	5	3	1	3	5	7	9	TA
LE	9	7	5	3	1	3	5	7	9	WI
TA	9	7	5	3	1	3	5	7	9	WI
IC	9	7	5	3	1	3	5	7	9	ST
IC	9	7	5	3	1	3	5	7	9	ER
ST	9	7	5	3	1	3	5	7	9	ER
WR	9	7	5	3	1	3	5	7	9	SU
WR	9	7	5	3	1	3	5	7	9	SC
WR	9	7	5	3	1	3	5	7	9	FR
WR	9	7	5	3	1	3	5	7	9	PR
SU	9	7	5	3	1	3	5	7	9	SC
SU	9	7	5	3	1	3	5	7	9	FR
SU	9	7	5	3	1	3	5	7	9	PR
SC	9	7	5	3	1	3	5	7	9	FR
SC	9	7	5	3	1	3	5	7	9	PR
FR	9	7	5	3	1	3	5	7	9	PR

acceptable. The same procedure is undertaken for the subcriteria in the other two criteria, i.e. economic importance and impact of climate change on the forest.

In the third step, the selected species must be prioritized in relation to each subcriterion. Then, the pairwise comparison and weighting procedure is undertaken. For the third criteria and for each of the five subcriteria, a prioritized list of species is obtained (see table 4). Once the eigenvectors for each subcriterion have been calculated with an acceptable level of consistency, the matrix of priority vectors for species and subcriteria can be constructed as shown in table 5. Figure 2 represents the factor's weight assigned to each subcriterion as a result of applying the eigenvector method in the pairwise comparison from the panel of experts. The priority matrix of the species for each subcriterion must be multiplied by the priority

matrix of this subcriterion in order to obtain an overall prioritized list of species. Figure 3 shows the optimal percentage distribution of the appropriate species to be used in the reforestation planning.


## 5. Conclusions

Climate change is not usually taken into account as a key factor in a specific stand of reforestation. Moreover, there is a lack of agreement on the weight of each species in a reforestation project, which is usually calculated in a discretionary way. Consequently, the success of a reforestation project is based mainly on experience learned by trial and error. Long-term tree regeneration demands somewhat more accuracy in the decision-making process. The proposed hybrid methodology with its consistency analysis can reduce



**Table 3.** Results from subcriteria pairwise comparisons using a 1–9 preference scale for each expert respect to the overall goal.

Pairwise criteria	Results for every expert											
	1st	2nd	3rd	4th	5th	6th	7th	8th	9th	10th	11th	12th
CB versus LE	5	7	1/5	9	5	1/3	9	9	9	9	5	1/7
CB versus TA	7	9	5	9	9	9	9	9	9	9	9	9
CB versus WI	5	7	1	9	9	9	9	9	9	9	9	9
LE versus TA	5	7	5	7	9	7	5	7	9	7	9	7
LE versus WI	5	5	3	7	5	5	7	9	7	1/3	7	5
TA versus WI	3	1/3	3	5	1	1/7	1/9	1/7	1/9	1/7	1/9	1/9
IC versus ST	3	1/3	5	1	1/3	1/3	1	1	1	1	1	1
IC versus ER	3	7	3	5	1/5	9	9	9	9	7	7	7
ST versus ER	3	9	7	7	3	9	9	7	7	7	9	9
WR versus SU	3	3	1	5	3	3	5	5	3	7	1/3	5
WR versus SC	1/3	1/5	1	1/5	1/3	1/5	1/5	1/3	1/7	1/5	1/5	3
WR versus FR	1/7	1/7	1/7	1/9	1/7	1/7	1/7	1/9	1/9	1/9	1/9	1
WR versus PR	1/5	1/3	1/5	1/7	1/3	1/7	1/5	1/7	1/7	1/9	1/7	3
SU versus SC	1/5	1/3	1/5	1	1/5	1/7	1	1/3	1/7	3	1/7	1/5
SU versus FR	1/9	1/7	1/9	1/5	1/7	1/9	1/7	1/7	1/9	1/9	1/3	1/9
SU versus PR	1/7	1/5	1/5	1/5	1/7	1/3	1/5	1/9	1/7	1/7	1/9	3
SC versus FR	1/3	1/3	1/7	1/5	1/5	1/3	1/5	1/5	1/5	1/3	1/5	1/7
SC versus PR	1/3	1/3	1/7	1	1/5	1/5	1/5	1/3	1	1/3	1/3	1/7
FR versus PR	3	1	5	3	3	3	5	3	7	5	1/7	7



random choices and integrate climate change criteria in the reforestation project.

Due to the large number of factors considered and the complexity of the system, reforestation planning requires decision makers to use multicriteria analyses in order to assess potential species for their suitability according to the criteria considered. Despite the variety of potential species for a reforestation project, the Delphi-AHP methodology described allows decision makers to focus on those providing a better suitability in each stand. At present, not only economic criteria are taken into account in reforestation but also those providing environmental services to society, always bearing in mind the criteria which might be related to

climate change, as promoted by the European Commission.

Not every factor in this study may have the same weight in the decision-making process. Therefore, the factors are assigned a weight based on the characteristics of the case studied, i.e. reforestation aimed at conserving biodiversity in the Mediterranean region. It is precisely in this assessment that the flexibility of this technique resides. As it can be observed in figure 2, the first column of each color shows the accumulated weight given for each subcriterion. The highest values correspond to FR (fire forest response) and PR (pests and diseases risk) because of the increasing importance of the impact of climate change. The

**Table 4.** Pairwise comparison matrix, eigenvector and consistency analysis for the six species with respect to each subcriteria of the ICC criteria.

RL_519886	Species	<i>Q. ilex</i>	<i>Q. faginea</i>	<i>F. ornus</i>	<i>P. spinosa</i>	<i>S. aria</i>	<i>C. monogyna</i>	Eigenvector
WR	<i>Q. ilex</i>	1.0000	8.6308	7.2994	8.6308	7.1481	5.4388	0.5426
	<i>Q. faginea</i>	0.1159	1.0000	1.0959	2.7375	0.3215	0.2306	0.0510
	<i>F. ornus</i>	0.1370	0.9125	1.0000	2.8566	0.3501	0.3355	0.0557
	<i>P. spinosa</i>	0.1159	0.3653	0.3501	1.0000	0.2059	0.1470	0.0278
	<i>S. aria</i>	0.1399	3.1105	2.8566	4.8568	1.0000	0.1866	0.1025
	<i>C. monogyna</i>	0.1839	4.3371	2.9808	6.8018	5.3603	1.0000	0.2203
$\lambda_{\max} = 6.5617$ CI = 0.1123 CR = 8.99% < 10% ( $n = 6$ )								
SU	<i>Q. ilex</i>	1.0000	6.7583	6.6183	4.9279	3.2666	1.2009	0.3331
	<i>Q. faginea</i>	0.1480	1.0000	4.7225	0.3236	0.1866	0.1159	0.0498
	<i>F. ornus</i>	0.1511	0.2118	1.0000	0.2178	0.1814	0.1380	0.0279
	<i>P. spinosa</i>	0.2029	3.0906	4.5919	1.0000	0.3676	0.1351	0.0816
	<i>S. aria</i>	0.3061	5.3603	5.5128	2.7200	1.0000	0.3501	0.1559
	<i>C. monogyna</i>	0.8327	8.6308	7.2478	7.4012	2.8566	1.0000	0.3517
$\lambda_{\max} = 6.5690$ CI = 0.1138 CR = 9.10% < 10% ( $n = 6$ )								
SC	<i>Q. ilex</i>	1.0000	1.3161	2.7200	2.6067	6.9014	6.7583	0.3346
	<i>Q. faginea</i>	0.7598	1.0000	2.8566	3.1305	6.8065	6.7583	0.3169
	<i>F. ornus</i>	0.3676	0.3501	1.0000	2.8566	2.8566	2.8566	0.1536
	<i>P. spinosa</i>	0.3836	0.3194	0.3501	1.0000	2.6067	3.1305	0.1050
	<i>S. aria</i>	0.1449	0.1469	0.3501	0.3836	1.0000	1.0959	0.0461
	<i>C. monogyna</i>	0.1480	0.1480	0.3501	0.3194	0.9125	1.0000	0.0438
$\lambda_{\max} = 6.1486$ CI = 0.0297 CR = 2.38% < 10% ( $n = 6$ )								
FR	<i>Q. ilex</i>	1.0000	0.1883	2.5736	4.6981	1.2211	6.5338	0.1901
	<i>Q. faginea</i>	5.3095	1.0000	7.2708	6.3862	1.2211	6.5846	0.4127
	<i>F. ornus</i>	0.3886	0.1375	1.0000	0.8189	0.1591	1.3493	0.0498
	<i>P. spinosa</i>	0.2129	0.1566	1.2211	1.0000	0.1643	0.3038	0.0403
	<i>S. aria</i>	0.8189	0.8189	6.2858	6.0874	1.0000	6.5846	0.2532
	<i>C. monogyna</i>	0.1530	0.1519	0.7411	3.2920	0.1519	1.0000	0.0539
$\lambda_{\max} = 6.5572$ CI = 0.1114 CR = 8.91% < 10% ( $n = 6$ )								
PR	<i>Q. ilex</i>	1.0000	4.7225	7.2994	2.8566	6.7583	1.0959	0.3091
	<i>Q. faginea</i>	0.2118	1.0000	4.7225	2.9808	2.7375	0.1380	0.1129
	<i>F. ornus</i>	0.1370	0.2118	1.0000	0.2934	0.3061	0.1159	0.0274
	<i>P. spinosa</i>	0.3501	0.3355	3.4087	1.0000	4.5919	0.1370	0.0883
	<i>S. aria</i>	0.1480	0.3653	3.2666	0.2178	1.0000	0.1159	0.0434
	<i>C. monogyna</i>	0.9125	7.2478	8.6308	7.2994	8.6308	1.0000	0.4190
$\lambda_{\max} = 6.6223$ CI = 0.1245 CR = 9.96% < 10% ( $n = 6$ )								

**Table 5.** Summary of the factor's priority and the species' weights for each subcriterion.

Criterion	Sub-criterion	Sub-criterion's weight	Selected species					
			<i>Q. ilex</i>	<i>Q. faginea</i>	<i>F. ornus</i>	<i>P. spinosa</i>	<i>S. aria</i>	<i>C. monogyna</i>
EES	CB	0.111	0.032	0.045	0.445	0.304	0.069	0.105
	LE	0.054	0.035	0.065	0.393	0.307	0.068	0.133
	TA	0.009	0.106	0.037	0.411	0.348	0.032	0.067
	WI	0.016	0.291	0.121	0.029	0.090	0.044	0.425
EIM	IC	0.072	0.031	0.045	0.278	0.453	0.082	0.111
	ST	0.083	0.032	0.031	0.296	0.459	0.071	0.112
	ER	0.014	0.237	0.111	0.038	0.542	0.034	0.037
ICC	WR	0.046	0.543	0.051	0.056	0.028	0.102	0.220
	SU	0.029	0.333	0.050	0.028	0.082	0.156	0.352
	SC	0.083	0.335	0.317	0.154	0.105	0.046	0.044
	FR	0.315	0.190	0.413	0.050	0.040	0.253	0.054
	PR	0.169	0.309	0.113	0.027	0.088	0.043	0.419




Figure 3. Optimum percentage distribution of the selected species for the reforestation.




Figure 4. Species assessment for each subcriterion proposed by panelists as a result of applying the AHP method.

subcriterion CB (conservation of biodiversity) is in third position in line with the aim of reforestation. As regards the economic criteria, experts have given more importance to those targeting the success of reforestation, like IC and ST.

The results in table 5 show that *F. ornus* and *Prunus spinosa* are the most suitable species for almost all subcriteria of the first criterion (0.4452 and 0.3044 respectively for CB, 0.3929 and 0.3069 for LE, 0.4107 and 0.3481 for TA), while for WI the highest values are for *C. monogyna* (0.4245) and *Quercus ilex* (0.2910). For the economic subcriteria the species considered best is again *P. spinosa* (0.4530 for IC, 0.4587 for ST and 0.5417 for ER), followed by *F. ornus* (0.2776 for IC and 0.2957 for ST). However, for ICC subcriteria *Q. ilex* is best placed (0.5426 for WR, 0.3331 for SU, 0.3346 for SC, 0.3091 for PR); the same happens with *Quercus faginea* (0.3169 for SC and 0.4127 for FR) and with *C. monogyna* (0.2203 for WR, 0.3517 for SU and 0.4190 for PR). Therefore, as can be observed in figure 3, the most suitable species for all the subcriteria studied are *Q. faginea* (19.75%) and *Q. ilex* (19.35%).

The assessment of the species selected under each criterion considered in the reforestation is shown in figure 4. The results demonstrate that some species, such as *Q. ilex*, *Q. faginea* or *P. spinosa*, offer a good balance between all the factors that might affect the success and viability of reforestation. Finally, the optimal percentage distribution of the species selected is highlighted in figure 3. Reforestation planning can be accomplished with several species according to the percentages achieved. This decision-support system provides a methodology to give support and coherence to the process of selecting species in a reforestation. In an increasingly complex world, we are forced every day to deal with complicated issues involving several variables or stakeholders that might interact. Reforestation planning is no exception. Moreover, there is a lack of knowledge on how to implement climate-change factors. The proposed hybrid methodology allows the joint analysis of both traditional and climate-change criteria in reforestation projects.

## References

- Akinci I and Akinci E 2010 Sustainable urban development: a case study on Goksu Delta *Conf. on Water Observation and Information System for Decision Support (Ohrid)* Balwois ([http://balwois.com/wp-content/uploads/old\\_proc/ffp-1366.pdf](http://balwois.com/wp-content/uploads/old_proc/ffp-1366.pdf))
- Amraoui M, Pereira M G, DaCamara C C and Calado T J 2015 Atmospheric conditions associated with extreme fire activity in the Western Mediterranean region *Sci. Total Environ.* **524**–525 32–9
- Barajas-Guzman M G and Barradas V L 2013 Costs and benefits of the use of mulches in deciduous tropical reforestation *Bot. Sci.* **91** 363–70
- Benito Garzon M, Alia R, Robson T M and Zavala M A 2011 Intra-specific variability and plasticity influence potential tree species distributions under climate change *Glob. Ecol. Biogeogr.* **20** 766–78
- Boucher Y, Grondin P and Auger I 2014 Land use history (1840–2005) and physiography as determinants of southern boreal forests *Landsc. Ecol.* **29** 437–50
- Buckley L B 2014 *Species Responses to Climate Change* (Oxford: Ecology-Oxford Bibliographies)
- Cabrera H M 2002 Ecophysiological responses of plants in ecosystems with Mediterranean-like climate and high mountain environments *Rev. Chil. Hist. Natl.* **75** 625–37 ([http://rchn.biologichile.cl/pdfs/2002/3/Cabrera\\_2002.pdf](http://rchn.biologichile.cl/pdfs/2002/3/Cabrera_2002.pdf))
- Canto-Perello J, Curiel-Esparza J and Calvo V 2013 Criticality and threat analysis on utility tunnels for planning security policies of utilities in urban underground space *Expert Syst. Appl.* **40** 4707–14
- Chen I C, Hill J K, Ohlemüller R, Roy D and Thomas C D 2011 Rapid range shifts of species associated with high levels of climate warming *Science* **333** 1024–6
- Collaborative Partnership on Forests (CPF) 2008 For a coordinated forest-sector response to climate change *Strategic framework for forests and climate change* (<http://fao.org/forestry/30137-08bfac75588a0dbd48f2af0ea63bcb4de.pdf>)
- Cram S, Sommer I, Morales L M, Oropeza O, Carmona E and Gonzalez-Medrano F 2006 Suitability of the vegetation types in Mexico's Tamaulipas state for the siting of hazardous waste treatment plan *J. Environ. Manage.* **80** 13–24
- Curiel-Esparza J and Canto-Perello J 2013 Selecting utilities placement techniques in urban underground engineering *Arch. Civ. Mech. Eng.* **13** 276–85
- Curiel-Esparza J, Cuenca-Ruiz M A, Martin-Utrillas M and Canto-Perello J 2014 Selecting a sustainable disinfection technique for wastewater reuse projects *Water* **6** 2732–47
- DeLucia E H, Nabity P D, Zavala J A and Berenbaum M R 2012 Climate change: resetting plant-insect interactions *Plant Physiol.* **160** 1677–85
- Denzel J K, Weidenhiller A and Golser M 2015 Property relationships between spruce long and structural timber *Scand. J. Forest Res.* **30** 617–23
- Diaz-Balteiro L and Romero C 2008 Making forestry decisions with multiple criteria: a review and an assessment *For. Ecol. Manage.* **255** 3222–41
- Dietz T and Jorgenson A K 2014 Towards a new view of sustainable development: human well-being and environmental stress *Environ. Res. Lett.* **9** 031001
- Dymov A A, Zhangurov E V and Hagedorn F 2015 Soil organic matter composition along altitudinal gradients in permafrost affected soils of the subpolar ural mountains *Catena* **131** 140–8
- Easterling D R, Meehl G A, Parmesan C, Changnon S A, Karl T R and Mearns L O 2000 Climate extremes: observations, modeling, and impacts *Science* **289** 2068–74
- Edwards D M, Jay M, Jensen F S, Lucas B, Marzano M, Montagne C, Peace A and Weiss G 2012a Public preferences across Europe for different forest stand types as sites for recreation *Ecol. Soc.* **17** 27
- Edwards D M, Jay M, Jensen F S, Lucas B, Marzano M, Montagne C, Peace A and Weiss G 2012b Public preferences for structural attributes of forests: towards a pan-European perspective *Forest Policy Econ.* **19** 12–9
- Espelta J M, Retana J and Habrouk A 2003 An economic and ecological multi-criteria evaluation of reforestation methods to recover burned *Pinus nigra* forests in NE Spain *For. Ecol. Manage.* **180** 185–98
- EUCOM European Commission 1998 *Communication from the Commission to the Council and to the European Parliament on a Forestry Strategy for the EU COM*(1998) 649 ([http://ec.europa.eu/agriculture/fore/publi/1998\\_649\\_en.pdf](http://ec.europa.eu/agriculture/fore/publi/1998_649_en.pdf))
- EUCOM European Commission 2006 *EU Forest Action Plan COM*(2006) 302 ([http://ec.europa.eu/agriculture/fore/action\\_plan/com\\_en.pdf](http://ec.europa.eu/agriculture/fore/action_plan/com_en.pdf))
- EUCOM European Commission 2011 *Communication from the Commission to the European Parliament, the Council, the Economic and Social Committee and the Committee of the Regions on an EU Biodiversity Strategy to 2020 COM*(2011)244 (<http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:52011DC0244&from=EN>)
- EUCOM European Commission 2013 *Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions on a new EU Forestry Strategy: for forests and the forest-based sector COM*(2013) 659 ([http://ec.europa.eu/agriculture/forest/strategy/staff-working-doc\\_en.pdf](http://ec.europa.eu/agriculture/forest/strategy/staff-working-doc_en.pdf))
- EUCON European Council 1999 *EU Forestry Strategy Council Resolution 1999/C/ 56/01* (<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:C:1999:056:0001:0004:EN:PDF>)
- EUPAR European Parliament 2011 *European Parliament Resolution of 11 May 2011 on the Commission Green paper on forest protection and information in the EU: preparing forests for climate change* (<http://www.europarl.europa.eu/sides/getDoc.do?type=TA&language=EN&reference=P7-TA-2011-226>)
- Ezebilo E E 2012 Forest stakeholder participation in improving game habitat in Swedish forests *Sustainability* **4** 1580–95
- FAO Food and Agriculture Organization of the United Nations 2013 Climate change guidelines for forest managers (<http://fao.org/docrep/018/i3383e/i3383e00.htm>)
- Fitter A H and Fitter R S R 2002 Rapid changes in flowering time in British plants *Science* **296** 1689–91
- Garcia-Lopez J M and Allue-Camacho C 2008 Phytoclimatic versatility and potential diversity of natural arboreal forest cover in peninsular Spain *Investigación Agraria: Sistemas y Recursos Forestales* **17** 297–307
- Garcia-Lopez J M and Allue-Camacho C 2010 Effects of climate change on the distribution of *Pinus sylvestris* L stands in Spain a phytoclimatic approach to defining management alternatives *For. Syst.* **19** 329–39
- Garson G D 2013 *The Delphi Method in Quantitative Research* (Asheboro: Statistical Associates Blue Book Series)
- Gilman S E, Urban M C, Tewksbury J, Gilchrist G W and Holt R D 2010 A framework for community interactions under climate change *Trends Ecol. Evol.* **25** 325–31
- Gomez-Aparicio L, Garcia-Valdes R, Ruiz-Benito P and Zavala M A 2011 Disentangling the relative importance of climate, size and competition on tree growth in Iberian forests: implications for forest management under global change *Glob. Change Biol.* **17** 2400–14
- Gomontean B, Gajasen J, Edwards-Jones G and Gajasen N 2008 The development of appropriate ecological criteria and indicators for community forest conservation using participatory methods: a case study in northeastern Thailand *Ecol. Indic.* **8** 614–24
- Gonzalez J R and Pukkala T 2007 Characterization of forest fires in Catalonia (north-east Spain) *Eur. J. For. Res.* **126** 421–9
- Hampel D, Janova J and Kadlec J 2015 Estimation of cost and revenue functions for reforestation system in dráhanska highlands *Int. Conf. on Numerical Analysis and Applied*



- Mathematics* (ICNAAM) AIP Conf. Proc. 1648 ed  
T E Simos and C Tsitouras (UNSP 720005) (doi:10.1063/1.4912942)
- Hannah L 2010 *Climate Change Biology* (Amsterdam: Elsevier)
- Hitsuma G, Morisawa T and Yagihashi T 2015 Orthotropic lateral branches contribute to shade tolerance and survival of *Thujaops dolabrata* var. *hondai* saplings by altering crown architecture and promoting layering *Botany* **93** 353–60
- Hockmann K, Tandy S, Lenz M, Reiser R, Conesa H M, Keller M, Studer B and Schulin R 2015 Antimony retention and release from drained and waterlogged shooting range soil under field conditions *Chemosphere* **134** 536–43
- Hofmann G E and Todgham A E 2010 Living in the now: physiological mechanisms to tolerate a rapidly changing environment *Annu. Rev. Physiol.* **72** 127–45
- Hunt D V L, Nash D and Rogers C D F 2014 Sustainable utility placement via multi-utility tunnels *Tunn Undergr Space Technol.* **39** 15–26
- Hóðar J A, Zamora R and Cayuela L 2012 Climate change and the incidence of a forest pest in Mediterranean ecosystems: can the North Atlantic Oscillation be used as a predictor? *Clim. Change* **113** 699–711
- IPCC Intergovernmental Panel of Climate Change 2014a *Climate Change 2014: Mitigation of Climate Change. Working Group III* ([http://report.mitigation2014.org/drafts/final-draft-postplenary/ipcc\\_wg3\\_ar5\\_final-draft\\_postplenary\\_chapter11.pdf](http://report.mitigation2014.org/drafts/final-draft-postplenary/ipcc_wg3_ar5_final-draft_postplenary_chapter11.pdf))
- IPCC Intergovernmental Panel of Climate Change 2014b *Climate Change 2014: Impacts, Adaptation and Vulnerability. Working Group II* ([http://ipcc-wg2.gov/AR5/images/uploads/WG2AR5\\_SPM\\_FINAL.pdf](http://ipcc-wg2.gov/AR5/images/uploads/WG2AR5_SPM_FINAL.pdf))
- IUCN International Union for Conservation of Nature 2004 *Afforestation and Reforestation for Climate Change Mitigation: Potentials for Pan-European Action* (<https://portals.iucn.org/library/efiles/documents/2004-066.pdf>)
- Kane B and Finn J T 2014 Factors affecting branch failures in open-grown trees during a snowstorm in Massachusetts, USA *Springer Plus* **3** 720
- Kaya T and Kahraman C 2011 Fuzzy multiple criteria forestry decision making based on an integrated VIKOR and AHP approach *Expert Syst. Appl.* **38** 7326–33
- Ki-moon B 2008 *United Nations—Reducing Emissions from Deforestation in Developing Countries (UN-REDD Programme) FAO's Forest and Climate Change Programme* (New York: United Nations Environment Programme)
- Kinoshita A M and Hogue T S 2015 Increased dry season water yield in burned watersheds in Southern California *Environ. Res. Lett.* **10** 014003
- Koprowski M and Dunker P 2012 Tree ring width and wood density as the indicators of climatic factors and insect outbreaks affecting spruce growth *Ecol. Indic.* **23** 332–7
- Kukavskaya E A, Buryak L V, Ivanova G A, Conard S G, Kalenskaya O P, Zhila S V and McRae D J 2013 Influence of logging on the effects of wildfire in Siberia *Environ. Res. Lett.* **8** 045034
- Lau J A and Lennon J T 2012 Rapid responses of soil microorganisms improve plant fitness in novel environments *Proc. Natl Acad. Sci. USA* **109** 14058–62
- Lenoir J, Gégout J C, Marquet P A, de Ruffray P and Brisse H 2008 A significant upward shift in plant species optimum elevation during the 20th century *Science* **320** 1768–71
- Lupp G, Konold W and Bastian O 2013 Landscape management and landscape changes towards more naturalness and wilderness: effects on scenic qualities—the case of the muritz national park in germany *J. Nat. Conserv.* **21** 10–21
- Marques A F, Ficko A, Kangas A, Rosset C, Ferreti F, Rasimaki J, Packalen T and Gordon S 2013 Empirical guidelines for forest management decision support systems based on the past experiences of the expert's community *For. Syst.* **22** 320–39
- Martin-Utrillas M, Azorin-Carrion A, Canto-Perello J and Curiel-Esparza J 2015a *Multi-Criteria Decision-Making Model for Establishing The Optimal Typology for Clinker Storage Silos* *Zkg Int.* **1-2** 50–8 ([http://www.zkg.de/en/artikel/zkg\\_Multi-criteria-decision-making\\_model\\_for\\_establishing\\_the\\_optimal\\_typology\\_2273460.html](http://www.zkg.de/en/artikel/zkg_Multi-criteria-decision-making_model_for_establishing_the_optimal_typology_2273460.html))
- Martin-Utrillas M, Juan-Garcia F, Canto-Perello J and Curiel-Esparza J 2015b Optimal infrastructure selection to boost regional sustainable economy *Int. J. Sustain. Dev. World Ecol.* **22** 30–8
- Martin-Utrillas M, Reyes-Medina M, Curiel-Esparza J and Canto-Perello J 2015c Hybrid method for selection of the optimal process of leachate treatment in waste treatment and valorization plants or landfills *Clean Technol. Environ. Policy.* **17** 873–85
- Mataix-Solera J, Cerda A, Arcenegui V, Jordan A and Zavala L M 2011 Fire effects on soil aggregation: a review *Earth-Sci. Rev.* **109** 44–60
- MCPFE Ministerial Conference on the Protection of Forests in Europe 2007 *Warsaw Ministerial Conference: Forests for quality of life. Pan-European Recommendations for Afforestation and Reforestation in the context of the UNFCCC, with a special focus on mitigation and adaptation to climate change* ([http://foresteurope.org/docs/other\\_meetings/2007/elm2/ELM\\_2\\_5\\_2007\\_Draft\\_Recommendations\\_on\\_Afforestation\\_and\\_Reforestation.pdf](http://foresteurope.org/docs/other_meetings/2007/elm2/ELM_2_5_2007_Draft_Recommendations_on_Afforestation_and_Reforestation.pdf))
- MCPFE Ministerial Conference on the Protection of Forests in Europe 2002 *Expert Level Meeting. Improved Pan-European Indicators for Sustainable Forest Management* ([http://foresteurope.org/docs/reporting/Vienna\\_Improved\\_Indicators.pdf](http://foresteurope.org/docs/reporting/Vienna_Improved_Indicators.pdf))
- MCPFE Ministerial Conference on the Protection of Forests in Europe 2011 *Oslo Ministerial Conference: European Forests 2020. Pan-European Guidelines for Afforestation and Reforestation with a special focus on the provisions of the UNFCCC* (<http://foresteurope.org/documentos/PanEuropeanAfforestationReforestationGuidelines.pdf>)
- Meddour-Sahar O, Meddour R, Leone V, Lovreglio R and Derridj A 2013 Analysis of forest fires causes and their motivations in northern Algeria: the Delphi method *Forest-Biogeoosci. Forestry* **6** 247–54
- Mendoza G A and Martins H 2006 Multi-criteria decision analysis in natural resource management: a critical review of methods and new modelling paradigms *For. Ecol. Manage.* **230** 1–22
- Moreno J M, Viedma O, Zavala G and Luna B 2011 Landscape variables influencing forest fires in central Spain *Int. J. Wildland Fire* **20** 678–89
- Navarro-Cerrillo R M, Griffith D M, Ramirez-Soria M J, Pariona W, Golicher D and Palacios G 2011 Enrichment of big-leaf mahogany (*Swietenia macrophylla* King) in logging gaps in Bolivia: The effects of planting method and silvicultural treatments on long-term seedling survival and growth *For. Ecol. Manage.* **262** 2271–80
- Newman J A, Anand M, Henry H A L, Hunt S and Gedalof Z 2011 *Climate Change Biology* (Cambridge, MA: CABI)
- Nielsen M R and Treue T 2012 Hunting for the benefits of joint forest management in the Eastern Afrotropical Biodiversity Hotspot: effects on Bushmeat hunters and wildlife in the Udzungwa Mountains *World Dev.* **40** 1224–39
- Ojea E, Ruiz-Benito P, Markandya A and Zavala M A 2012 Wood provisioning in Mediterranean forests: a bottom-up spatial valuation approach *Forest Policy Econ.* **20** 78–88
- Orsi F, Geneletti D and Newton A C 2011 Towards a common set of criteria and indicators to identify forest restoration priorities: an expert panel-based approach *Ecol. Indic.* **11** 337–47
- Paoletti E 2005 Ozone slows stomatal response to light and leaf wounding in a Mediterranean evergreen broadleaf *Arbutus unedo*. *Environ. Pollut.* **134** 439–45
- Parmesan C 2006 Ecological and evolutionary responses to recent climate change *Annu. Rev. Ecol. Evol. Systemat.* **37** 637–69
- PATFOR 2013 *Territorial Forestry Action Plan from Comunitat Valenciana. Conselleria de Infraestructuras, Territorio y Medio Ambiente (CMA) Decreto 58/2013, de 3 de mayo, del Consell, por el que se aprueba el Plan de Accion Territorial Forestal de la*



- Comunitat Valenciana (PATFOR) DOGV 7019 de 08052013 ([http://docv.gva.es/portal/ficha\\_disposicion\\_pc.jsp?sig=004345/2013&L=1](http://docv.gva.es/portal/ficha_disposicion_pc.jsp?sig=004345/2013&L=1))
- Peman Garcia J, Navarro Cerrillo R M and Serrada Hierro R 2006 Eleccion de especies en las repoblaciones forestales *Investigaciones Agrarias: Sistemas de Recursos Forestales* **15** 87–102 ([http://www.inia.es/gcontrec/pub/087-102-\(29\)-Eleccion\\_1169111282265.pdf](http://www.inia.es/gcontrec/pub/087-102-(29)-Eleccion_1169111282265.pdf))
- Pettorelli N 2012 Climate change as a main driver of ecological research *J. Appl. Ecol.* **49** 542–5
- Ramos M C and Martinez-Casasnovas J A 2015 Climate change influence on runoff and soil losses in a rainfed basin with Mediterranean climate *Nat. Hazards* **78** 1065–89
- Reilly J R and Elderd B D 2014 Effects of biological control on long-term population dynamics: identifying unexpected outcomes *J. Appl. Ecol.* **51** 90–101
- Roura-Pascual N *et al* 2009 Ecology and management of alien plant invasions in South African fynbos: accommodating key complexities in objective decision making *Biol. Conserv.* **142** 1595–604
- Ruiz-Benito P, Lines E R, Gomez-Aparicio L, Zavala M A and Coomes D A 2013 Patterns and drivers of tree mortality in Iberian forests: climatic effects are modified by competition *PLoS ONE* **8** e56843
- Saaty T L 2008 Decision making with the analytic hierarchy process *Int. J. Serv. Sci.* **1** 83–98
- Saaty T L 2012 *Decision Making for Leaders The Analytic Hierarchy Process for Decisions in a Complex World* (Pittsburgh: Rws Publications)
- Saaty T L 2013 The modern science of Multicriteria Decision Making and its practical applications: the AHP/ANP approach *Oper. Res.* **61** 1101–18
- Salvini G, Herold M, De Sy V, Kissinger G, Brockhaus M and Skutsch M 2014 How countries link REDD+ interventions to drivers in their readiness plans: implications for monitoring systems *Environ. Res. Lett.* **9** 074004
- Sanchez-Gomez D, Valladares F and Zavala M A 2006 Performance of seedlings of Mediterranean woody species under experimental gradients of irradiance and water availability: trade-offs and evidence for niche differentiation *New Phytol.* **170** 795–805
- Sanchez-Salguero R, Navarro-Cerrillo R M, Swetnam T W and Zavala M A 2012 Is drought the main decline factor at the rear edge of Europe? The case of southern Iberian pine plantations *For. Ecol. Manage.* **271** 158–69
- Schröter D *et al* 2005 Ecosystem service supply and vulnerability to global change in Europe *Science* **310** 1333–7
- Selkima M, Gonzalez-Olabarria J R and Pukkala T 2012 Site and stand characteristics related to surface erosion occurrence in forests of Catalonia (Spain) *Eur. J. For. Res.* **131** 727–38
- Sexton J P, McIntyre P J, Angert A L and Rice K J 2009 Evolution and ecology of species range limits *Annu. Rev. Ecol. Evol. Systemat.* **40** 415–36
- Sullivan P F, Ellison S B Z, McNown R W, Brownlee A H and Sveinbjornsson B 2015 Evidence of soil nutrient availability as the proximate constraint on growth of treeline trees in northwest alaska *Ecol.* **96** 716–27
- Urbietta I R, Perez-Ramos I M, Zavala M A, Maranon T and Kobe R K 2008 Soil water content and emergence time control seedling establishment in three co-occurring Mediterranean oak species *Can. J. Forest Res.* **38** 2382–93
- Urli M, Delzon S, Eyermann A, Couallier V, Garcia-Valdes R, Zavala M A and Porte A J 2014 Inferring shifts in tree species distribution using asymmetric distribution curves: a case study in the Iberian mountains *J. Veg. Sci.* **25** 147–59
- Vautard R *et al* 2014 The European climate under a 2 °C global warming *Environ. Res. Lett.* **9** 034006
- Vazquez A, Gonzalez-Alonso F, Cuevas J M, Calle A and Casanova J L 2002 Relationship between drought indicators based on remote sensing and forest fires incidence in Spain *Observing our Environment for Space: New Solutions for a new Millennium 21st Annual Symp. of the European-Association-of-Remote-Sensing-Laboratories 111 + ed G Begni (Paris)*
- Velmurugan A, Swarnam T P and Lal R 2015 Effect of land shaping on soil properties and crop yield in tsunami inundated coastal soils of Southern Andaman Island *Agric. Ecosyst. Environ.* **206** 1–9
- Vicente-Serrano S M *et al* 2014 Evidence of increasing drought severity caused by temperature in rise southern Europe *Environ. Res. Lett.* **9** 044001
- Williams S E, Shoo L P, Isaac J L, Hoffmann A A and Langham G 2008 Towards an integrated framework for assessing the vulnerability of species to climate change *PLoS Biol.* **6** e325
- Xu Z L *et al* 2013 Estimating realized and potential carbon storage benefits from reforestation and afforestation under climate change: a case study of the Qinghai spruce forests in the Qilian Mountains, northwestern China *Mitig. Adapt. Strateg. Glob. Chang.* **18** 1257–68
- Zhou S, Yin Y, Xu W, Ji Z, Caldwell I and Ren J 2007 The costs and benefits of reforestation in Liping County, Guizhou Province, China *J. Environ. Manage.* **85** 722–35