

## Towards typologies of urban climate and global environmental change

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 Environ. Res. Lett. 10 101001

(<http://iopscience.iop.org/1748-9326/10/10/101001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.105

This content was downloaded on 13/04/2017 at 08:27

Please note that [terms and conditions apply](#).

You may also be interested in:

[Impact of urbanization on US surface climate](#)

Lahouari Bounoua, Ping Zhang, Georgy Mostovoy et al.

[Investigating the climate impacts of urbanization and the potential for cool roofs to counter future climate change in Southern California](#)

P Vahmani, F Sun, A Hall et al.

[Interactions between urban vegetation and surface urban heat islands: a case study in the Boston metropolitan region](#)

Eli K Melaas, Jonathan A Wang, David L Miller et al.

[Green and cool roofs to mitigate urban heat island effects in the Chicago metropolitan area: evaluation with a regional climate model](#)

A Sharma, P Conry, H J S Fernando et al.

[Contrasting responses of urban and rural surface energy budgets to heat waves explain synergies between urban heat islands and heat waves](#)

Dan Li, Ting Sun, Maofeng Liu et al.

[Diurnal changes in urban boundary layer environment induced by urban greening](#)

Jiyun Song and Zhi-Hua Wang

[Contrasting impacts of urban forms on the future thermal environment: example of Beijing metropolitan area](#)

Long Yang, Dev Niyogi, Mukul Tewari et al.

[The effectiveness of cool and green roofs as urban heat island mitigation strategies](#)

Dan Li, Elie Bou-Zeid and Michael Oppenheimer

## Environmental Research Letters



## PERSPECTIVE

## OPEN ACCESS

PUBLISHED  
14 October 2015

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



## Towards typologies of urban climate and global environmental change

Felix Creutzig

Mercator Research Institute on Global Commons and Climate Change, Torgauer Straße 12-15, 10829 Berlin, Germany

**Keywords:** urban climate, surface urban heat island, climate adaptation, climate mitigation, typology**Abstract**

The beauty of cities is that every city is different. From the homogenizing perspective of global environmental change that speaks trouble. We need an understanding of which kind of cities can contribute what kind of measures to mitigate and adapt to global environmental change. Typologies of cities offer a bridge between the idiosyncratic and the global. Bounoua *et al* (2015 *Environ. Res. Lett.* **10** 084010) analyse the impact of urbanization on surface climate. We discuss their results and suggest avenues for further systematic analysis.

Urbanization represents a comparably small land-use take globally, but the change in land use intensifies, with infrastructures built to stay. This raises decisive questions on the change in urban climate, the modulation of global warming at local scale, and the change in vegetation at different spatial scales. Bounoua *et al* (2015) contribute important insights to this blooming research field by investigating the surface urban heat island (SUHI) effect in urban regions in the United States. The authors use two different data sets with different spatial resolution: one covers urban land cover at 30 m resolution (Land-sat-based ISA), and the other biophysical products at 500 m resolution at 8 day-intervals (MODIS). Relying on meteorological data as well as heat absorption functions Bounoua, Zhang *et al* model surface temperature of ten US cities, representing different ecoregions. By calculating the difference between impervious surface area and the vegetated land surrounding them, they calculate the SUHI of these cities.

The study contains insights that are of interest to the wider scientific community. The paper demonstrates the specific influence of vegetation class—which determines to large degree the temperature differences between built environment and surrounding area. As a result, the SUHI varies between 3.3 °C (broadleaf deciduous forest, Washington, DC) and 2.2 °C–2.3 °C (temperature grassland, Chicago), and a negative SUHI of –2.5 °C (desert, Phoenix). This makes intuitive sense, as broadleaf trees offer more cooling by transpiration, in contrast to arid desert areas. The SUHI is also more pronounced where

absolute temperatures are lower (e.g. comparing Washington DC with Atlanta). The reason is that higher temperatures lead to shutdown of plant transpiration (or more formally: stomatal closure), by this ceasing their cooling function.

These data are valuable for further research aiming to identify appropriate mitigation and adaptation strategies for different classes of cities. The aim is to build action-oriented typologies of cities.

Hence, it is valuable to reflect the results of Bounoua *et al* in the context of mitigating urban heat islands. In a study on urban adaptation in the US, (Georgescu *et al* 2014) report that cool highly reflective white roofs are a more effective cooling strategy than adding vegetation to roofs. But the difference between cool roofing and green roofing is more accentuated in arid states like California (1.2 °C additional cooling) than in humid states like Florida (0.2 °C additional cooling). Together these studies suggest tentative mitigation portfolios (Georgescu *et al* 2015) for different types of urban climates:

- especially but not only in arid environments: increase the reflectivity of the built environment;
- if space and water is available: plant urban trees, possibly heat resistant types. Reduce impervious surface where possible, e.g. modify parking spaces;
- especially if space is limited: Include green roofs as part of portfolios.

Such a typology would also need to systematically include a number of confounding factors (Georgescu *et al* 2015). For example, reflectivity increases would be compromised by seasonal sandstorms; reflectivity increases might also impact hydrological cycles by reducing precipitation at regional scale. In addition, water scarcity might induce detrimental trade-offs on urban scale or beyond. A total weighting of mitigation typologies will not only depend on the expected change in extreme heat events for cities, but also demographic change, and hence total exposure (Jones *et al* 2015).

Clearly, these strategies also influence climate mitigation strategies. For example, cool roofs or increased urban vegetation reduce cooling energy demand, but in the case of reflectivity modifications also can increase urban heating demand in winter. In turn, adding green spaces is likely to induce higher health perception and significantly less cardio-metabolic conditions (Kardan *et al* 2015). Hence, the wider goal is the integration of urban mitigation and adaptation strategies, and possibly even urban quality of life, into comprehensive typologies, as called for by Solecki *et al* (2015). Such an comprehensive endeavor would have energy-use and GHG emission typologies of cities as its second pillar, in addition to the pillar of an urban climate typology. In fact, our own typologies, relying on hierarchical tree regression, clearly demonstrate that local climate (heating degree days and cooling degree days) is an important discriminator of cities on global level (Creutzig *et al* 2015) and on local human settlement level (e.g., in England: Baiocchi *et al* 2015). In these city typologies, other discriminators include urban form, economic well-being, and fuel prices.

A second interesting result of Bounoua *et al* addresses the issue of land carbon storage in urban areas and could also become element of the suggested comprehensive typologies. Bounoua *et al* calculate the counterfactual stored in urban areas by extrapolating the carbon content of the surrounding vegetation. They find that impervious surface—representing 1.1% of the total continental US land—replaces 0.9%–1.8% of the total US land carbon uptake, in line with (Imhoff *et al* 2004). This is a surprisingly high effect, given that agricultural land displays a much lower relative change in carbon (plus 5.0% representing 32.1% of total continental US land).

Such results are important as global model of net primary productivity remain ignorant of the urban contribution, and model urban carbon uptake with fixed parameters, a situation that modellers are eager to change (Haberl *et al* 2007). But of course, the result is also significant in its own right: reducing the impervious surface in cities is likely to make a measurable contribution to increasing total land carbon uptake.

There are two other but related concerns that point into fruitful further research directions: geographical scope, and integration with theory. First, the systematic analysis of urban heat islands is currently US centric. As other world regions display different climates, and often very different urban forms, analysis of other world regions would likely enrich any typology of urban climate and global environmental change. Second, urban economics provides a rich potential to relate urban dynamics to empirical insights on environmental and climate change outcomes (e.g., Viguié and Hallegatte 2012, Creutzig 2014). It would be particularly interesting to scrutinize the role of urban form and population density and to elucidate what how different mitigation and adaptation constraints interact to produce ‘sustainability windows of urban form’ (Lohrey and Creutzig 2015).

## References

- Baiocchi G, Creutzig F, Minx J and Pichler P-P 2015 A spatial typology of human settlements and their CO<sub>2</sub> emissions in England *Glob. Environ. Change* **34** 13–21
- Bounoua L, Zhang P, Mostovoy G, Thome K, Masek J, Imhoff M, Shepherd M, Quattrochi D, Santanello J and Silva J 2015 Impact of urbanization on US surface climate *Environ. Res. Lett.* **10** 084010
- Creutzig F 2014 How fuel prices determine public transport infrastructure, modal shares and urban form *Urban Clim.* **10** 63–76
- Creutzig F, Baiocchi G, Bierkandt R, Pichler P-P and Seto K C 2015 Global typology of urban energy use and potentials for an urbanization mitigation wedge *Proc. Natl Acad. Sci. USA* **112** 6283–8
- Georgescu M, Chow W, Wang Z, Brazel A, Trapido-Lurie B, Roth M and Benson-Lira V 2015 Prioritizing urban sustainability solutions: coordinated approaches must incorporate scale-dependent built environment induced effects *Environ. Res. Lett.* **10** 061001
- Georgescu M, Morefield P E, Bierwagen B G and Weaver C P 2014 Urban adaptation can roll back warming of emerging megapolitan regions *Proc. Natl Acad. Sci. USA* **111** 2909–14
- Haberl H, Erb K-H, Krausmann F, Gaube V, Bondeau A, Plutzer C, Gingrich S, Lucht W and Fischer-Kowalski M 2007 Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems *Proc. Natl Acad. Sci. USA* **104** 12942–7
- Imhoff M L, Bounoua L, DeFries R, Lawrence W T, Stutzer D, Tucker C J and Ricketts T 2004 The consequences of urban land transformation on net primary productivity in the United States *Remote Sens. Environ.* **89** 434–43
- Jones B, O'Neill B C, McDaniel L, McGinnis S, Mearns L O and Tebaldi C 2015 Future population exposure to US heat extremes *Nat. Clim. Change* **5** 652–5
- Kardan O, Gozdyra P, Misic B, Moola F, Palmer L J, Paus T and Berman M G 2015 Neighborhood greenspace and health in a large urban center *Sci. Rep.* **5** 11610
- Lohrey S and Creutzig F 2015 A 'sustainability window' of urban form *Transp. Res. D* in press
- Solecki W *et al* 2015 A conceptual framework for an urban areas typology to integrate climate change mitigation and adaptation *Urban Clim.* in press
- Viguié V and Hallegatte S 2012 Trade-offs and synergies in urban climate policies *Nat. Clim. Change* **2** 334–7